

Table: Annual Average OMI Tropospheric NO₂ VCD in 2005 and 2013, Total Reduction (%) for 2005-2013, and Annual Trend (%/yr) for 20 U.S. Metropolitan Areas.

Cities	Tropospheric NO ₂ VCD (10 ¹⁵ molec. cm ⁻²)		2005-2013 reduction (%)	Annual trend (%/year)	
	2005	2013		2005-2008	2010-2013
New York	16.5	11.2	40.9	-6.7	1.3
Los Angeles	20.5	11.8	43.6	-4.3	-4.3
Chicago	10.8	8.8	29.7	-2.6	-1.0
Dallas	5.7	3.8	38.5	-4.1	-2.0
Houston	8.5	5.2	39.4	-7.9	0.9
Philadelphia	10.9	6.7	47.1	-5.2	-0.4
Washington DC	9.6	4.7	48.6	-6.7	-3.4
Miami	4.3	3.3	23.0	-6.4	0.5
Atlanta	9.2	4.5	46.6	-7.9	-2.2
Boston	7.1	5.0	35.7	-8.1	-0.3
San Francisco	5.0	4.0	30.2	-2.3	-1.8
Phoenix	6.8	4.0	46.6	-6.7	-0.8
Detroit	9.4	5.2	25.1	-3.1	-1.3
Seattle	5.7	3.9	25.5	-2.9	-0.6
Minneapolis	4.9	3.1	23.4	-2.8	-1.9
San Diego	5.8	3.9	36.2	-5.0	-3.3
Tampa	5.0	2.8	47.0	-6.6	0.2
St. Louis	5.8	3.9	31.5	-5.0	0.6
Baltimore	8.6	5.4	40.2	-5.5	-1.5
Denver	7.0	5.6	26.3	-4.9	-2.8